



Buckeye®

SANICARE TBX™

DETERGENT DISINFECTANT PUMP SPRAY



SANICARE TBX, a ready-to-use, quaternary ammonium chloride based product, is a general non-acid cleaner and disinfectant for use in hospitals, nursing homes, schools, food service and lodging establishments, retail businesses and athlete/recreational facilities. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces such as floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, bathrooms, shower stalls, bathtubs and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** May be used in the kitchen on counters, sinks, appliances and stove tops. A rinse with potable water is required for surfaces in direct contact with food. In addition, **SANICARE TBX** deodorizes those areas which are generally hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.



This product exhibits disinfectant efficacy against Mycobacterium tuberculosis BCG (Mycobacterium bovis) at 20 degrees Centigrade when the treated surface is allowed to remain wet for **5 minutes**.

SANICARE TBX, when used on environmental, inanimate hard surfaces, exhibits virucidal activity against Hepatitis A Virus (HAV), Poliovirus Type 1, Canine Parvovirus, **Hepatitis B Virus (HBV)**, **Hepatitis C Virus (HCV)**, **Bovine Viral Diarrhea Virus (BVDV)**, **Human Coronavirus, SARS associated Coronavirus, Avian Influenza A strain H3N2, Avian Influenza A strain H9N2, Paramyxovirus (Mumps), Rhinovirus type 39, Rotavirus, HIV-1 (associated with AIDS), Norovirus (Norwalk virus), Feline Calicivirus, and Rabies Virus.** See product label for contact times.

SANICARE TBX kills the following organisms on hard non-porous inanimate surfaces:

BACTERICIDAL ACTIVITY: Staphylococcus aureus, Salmonella (choleraesuis) enterica, Escherichia coli, Pseudomonas aeruginosa, Escherichia coli 0157:H7, Listeria monocytogenes, Yersinia enterocolitica, Enterococcus faecium, Corynebacterium ammoniagenes, Salmonella (typhi) enterica, and Streptococcus pyogenes.

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY: Methicillin resistant Staphylococcus aureus (MRSA); Vancomycin intermediate resistant Staphylococcus aureus (VISA); Vancomycin resistant Enterococcus faecalis (VRE); Methicillin resistant Staphylococcus epidermidis (MRSE); Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA), (NRS 123) Genotype USA400; Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA), (NRS 384) Genotype USA300.

SANICARE TBX is fungicidal against the pathogenic fungus, Trichophyton mentagrophytes (Athlete's Foot Fungus)(cause of ringworm), when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet. To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.



SANICARE TBX™

DETERGENT DISINFECTANT PUMP SPRAY

DIRECTIONS & TECHNICAL SPECIFICATIONS

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DISINFECTION, DEODORIZING AND CLEANING

Remove gross filth or heavy soil prior to application of the product. Hold container 6 to 8 inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for Bloodborne Pathogens: allow surfaces to remain wet for HIV-1 **1 minute** and for HBV and HCV **5 minutes**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

One-step cleaning and disinfection of toilet bowls and urinals: Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of **SANICARE TBX** evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 **1 minute** and for HBV and HCV **5 minutes**) and flush.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

EPA REG. NO 1839-83-559
DIN: 02258226

EPA EST. NO 559-MO-1
U.S. Patent No. 5,444,094

Available in:



TECHNICAL SPECIFICATIONS

Specific Gravity	1.01
pH	11.7 ± 0.2
Color	Pale Clear Yellow
Odor	Lemon
Freeze-Thaw Stability	Unaffected

Buckeye International, Inc.

2700 Wagner Place • Maryland Heights • Missouri 63043 • 800.321.2583
www.buckeyeinternational.com